

**Appendix I (a)**  
**Human Surveillance Case Definition**  
(Revised July 4, 2005)

**Section A: Case Definitions**

The current Case Definitions were drafted with available information at the time of writing. Case Definitions and Diagnostic Test Criteria are subject to change as new information becomes available.

**1) West Nile Virus Neurological Syndrome (WNNS):**

**Clinical Criteria:**

History of exposure in an area where WN virus (WNV) activity is occurring<sup>1</sup>

**OR**

history of exposure to an alternative mode of transmission<sup>2</sup>

**AND**

onset of fever

**AND NEW ONSET OF AT LEAST ONE of the following:**

- encephalitis (acute signs of central or peripheral neurologic dysfunction),  
or
- viral meningitis (pleocytosis and signs of infection e.g. headache, nuchal rigidity), or
- acute flaccid paralysis (e.g. poliomyelitis-like syndrome or Guillain-Barré-like syndrome)<sup>3</sup> or
- movement disorders (e.g., tremor, myoclonus) or
- Parkinsonism or Parkinsonia like conditions (e.g., cogwheel rigidity, bradykinesia, postural instability) or
- other neurological syndromes as defined in the note below

<sup>1</sup>History of exposure when and where West Nile virus transmission is present, or could be present, or history of travel to an area with confirmed WNV activity in birds, horses, other mammals, sentinel chickens, mosquitoes, or humans.

<sup>2</sup>Alternative modes of transmission, identified to date, include: laboratory-acquired; *in utero*; receipt of blood components; organ/tissue transplant; and, possibly via breast milk.

<sup>3</sup> A person with WNV-associated acute flaccid paralysis may present with or without fever or mental status changes. Altered mental status could range from confusion to coma with or without additional signs of brain dysfunction (e.g. paralysis, cranial nerve palsies, sensory deficits, abnormal reflexes, generalized convulsions and abnormal movements). Acute flaccid paralysis with respiratory failure is also a problem.

**Note:** A significant feature of West Nile viral neurologic illness may be marked muscle weakness that is more frequently unilateral, but could be bilateral. WNV should be considered in the differential diagnosis of all suspected cases of acute flaccid paralysis with or without sensory deficit. WNV-associated weakness typically affects one or more limbs (sometimes affecting one limb only). Muscle weakness may be the sole presenting feature of WNV illness (in the absence of other

neurologic features) or may develop in the setting of fever, altered reflexes, meningitis or encephalitis. Weakness typically develops early in the course of clinical infection. Patients should be carefully monitored for evolving weakness and in particular for acute neuromuscular respiratory failure, which is a severe manifestation associated with high morbidity and mortality.

**For the purpose of WNV Neurological Syndrome Classification, muscle weakness is characterized by severe (Polio-like), non-transient and prolonged symptoms.**

Electromyography (EMG) and lumbar puncture should be performed to differentiate WNV paralysis from the acute demyelinating polyneuropathy (Guillain-Barré syndrome). Lymphocytic pleocytosis (an increase in WBC with a predominance of lymphocytes in the cerebrospinal fluid [CSF]) is commonly seen in acute flaccid paralysis due to WNV.

Other emerging clinical syndromes, identified during 2002 included, but were not limited to the following: myelopathy, rhabdomyolysis (acute destruction of skeletal muscle cells), peripheral neuropathy; polyradiculoneuropathy; optic neuritis; and acute demyelinating encephalomyelitis (ADEM). Ophthalmologic conditions including chorioretinitis and vitritis were also reported. Facial weakness was also reported. Myocarditis, pancreatitis and fulminant hepatitis have not been identified in North America, but were reported in outbreaks of WNV in South Africa. "Aseptic" meningitis without encephalitis or flaccid paralysis occurring in August and September when WNV is circulating may be due to non-polio enteroviruses circulating at the same time. This should be considered in the differential diagnosis.

[Sejvar J et al. *JAMA* (2003) Vol.290 (4) p. 511-515, Sejvar, J. et al. *Emerg Infect Dis* (2003) Vol 9 (7) p.788-93 and Burton, JM et al *Can. J. Neurol. Sci.* (2004) Vol.31 (2) p.185-193]

### **Suspect WN Neurological Syndrome Case:**

Clinical criteria IN THE ABSENCE OF OR PENDING diagnostic test criteria (see below) AND IN THE ABSENCE of any other obvious cause.

### **Probable WN Neurological Syndrome Case:**

Clinical criteria AND AT LEAST ONE of the probable case diagnostic test criteria (see below).

### **Confirmed WN Neurological Syndrome Case:**

Clinical criteria AND AT LEAST ONE of the confirmed case diagnostic test criteria (see below).

## **2) West Nile Virus Non-Neurological Syndrome (WN Non-NS):**

### **Clinical Criteria:**

History of exposure in an area where WN virus (WNV) activity is occurring<sup>1</sup>

**OR**

History of exposure to an alternative mode of transmission<sup>2</sup>

**AND AT LEAST TWO** of the following<sup>5</sup>:

- fever,
- myalgia<sup>6</sup>,
- arthralgia,
- headache,
- fatigue,
- lymphadenopathy,
- maculopapular rash

<sup>1</sup>History of exposure when and where West Nile virus transmission is present, or could be present, or history of travel to an area with confirmed WNV activity in birds, horses, other mammals, sentinel chickens, mosquitoes, or humans.

<sup>2</sup>Alternative modes of transmission, identified to date, include: laboratory-acquired; *in utero*; receipt of blood components; organ/tissue transplant; and, possibly via breast milk.

<sup>5</sup>It is possible that other clinical signs and symptoms could be identified that have not been listed and may accompany probable case or confirmed case diagnostic test criteria. For example, gastrointestinal (GI) symptoms were seen in many WNV patients in Canada and the USA in 2003 and 2004.

<sup>6</sup> Muscle weakness may be a presenting feature of WNV illness. **For the purpose of WNV Non-Neurological Syndrome classification, muscle weakness or myalgia (muscle aches and pains) is characterized by mild, transient, unlikely prolonged symptoms that are not caused by motor neuropathy.**

**Suspect WN Non-Neurological Syndrome Case:**

Clinical criteria IN THE ABSENCE OF OR PENDING diagnostic test criteria (see below) AND IN THE ABSENCE of any other obvious cause.

**Probable WN Non-Neurological Syndrome Case:**

Clinical criteria AND AT LEAST ONE of the probable case diagnostic test criteria (see below)

**Confirmed WN Non-Neurological Syndrome Case:**

Clinical criteria AND AT LEAST ONE of the confirmed case diagnostic test criteria (see below)

**3) West Nile Virus Asymptomatic Infection (WNAI)<sup>7</sup>:**

**Probable WN Asymptomatic Infection Case:**

Probable case diagnostic test criteria (see below) IN THE ABSENCE of clinical criteria

**Confirmed WN Asymptomatic Infection Case:**

Confirmed case diagnostic test criteria (see below) IN THE ABSENCE of clinical criteria

<sup>7</sup>This category could include asymptomatic blood donors whose blood is screened using a Nucleic Acid Amplification Test (NAT), by Blood Operators (i.e. Canadian Blood Services or Hema-Quebec) and is subsequently brought to the attention of public health officials. The NAT that will be used by Blood Operators in Canada is designed to detect all viruses in the Japanese encephalitis (JE) serocomplex. The JE serocomplex includes WN virus and 9 other viruses, although from this group only WN virus and St Louis encephalitis virus are currently endemic to parts of North America. Blood Operators in Canada perform a supplementary WN virus-specific NAT following any positive donor screen test result.

## **Section B: West Nile Virus Diagnostic Test Criteria:**

### **Probable Case Diagnostic Test Criteria:**

#### **AT LEAST ONE of the following:**

Detection of flavivirus antibodies in a single serum or CSF sample using a WN virus IgM ELISA <sup>8</sup> without confirmatory neutralization serology (e.g. Plaque Reduction Neutralization Test [PRNT]) <b>OR</b>
A 4-fold or greater change in flavivirus HI titres in paired acute and convalescent sera or demonstration of a seroconversion using a WN virus IgG ELISA <sup>8</sup> <b>OR</b>
A titre of $\geq 1:320$ in a single WN virus HI test, or an elevated titre in a WN virus IgG ELISA, with a confirmatory PRNT result <b>OR</b> [Note: A confirmatory PRNT or other kind of neutralization assay is not required in a health jurisdiction/authority where cases have already been confirmed in the current year]
Demonstration of Japanese encephalitis (JE) serocomplex-specific genomic sequences in blood by NAT screening on donor blood, by Blood Operators in Canada.

<sup>8</sup> Both CDC and commercial IgM / IgG ELISAs are now available for front line serological testing. Refer to appropriate assay procedures and kit inserts for the interpretation of test results.

**Note:** WNV IgM antibody may persist for more than a year and the demonstration of IgM antibodies in a patient's serum, particularly in residents of endemic areas, may not be diagnostic of an *acute* WN viral infection. Seroconversion (by HI, IgG ELISA or PRNT assays) demonstrates a current WNV infection. Therefore, the collection of acute and convalescent sera for serologic analysis is particularly important to rule out diagnostic misinterpretation early in the WNV season (e.g. May, June) and to identify initial cases in a specific jurisdiction. However, it should be noted that seroconversions may not always be documented due to timing of acute sample collection (i.e. titres in acute sera may have already peaked). If static titres are observed in acute and convalescent paired sera, it is still possible the case may represent a recent infection. To help resolve this, the use of IgG avidity testing <sup>9</sup> may be considered to distinguish between current and past infection. The presence of both IgM antibody and low avidity IgG in a patient's convalescent serum sample are consistent with current cases of viral associated illness. However test results that show the presence of IgM and high avidity IgG are indicative of exposures that have occurred in the previous season. Immunocompromised individuals may not be able to mount an immune response necessary for a serological diagnosis. West Nile virus diagnostic test criteria for these individuals should be discussed with a medical microbiologist.

<sup>9</sup> Early in infection the immune system generates antibodies that bind relatively weakly to viral antigen (low avidity). As the infection proceeds, an increasing percentage of newly generated IgG antibody displays higher binding affinity to virus antigen and thus avidity also rises (Note: avidity is usually measured based upon the ability of IgG to dissociate from antigen preparations)

after incubation with a solution of urea). As long as high avidity IgG is not yet detected in the serum it can be assumed that the individual was exposed to the viral agent during a recent exposure. With respect to WNV infection it has not been precisely determined when (i.e. post-exposure) high avidity antibodies reach levels in serum that can be accurately detected by serological assays (there may be significant variation depending on the individual). However, it has been shown that greater than 95% of sera collected from individuals exposed to WNV 6-8 months previously will have IgG antibodies that bind strongly to viral antigen and will give high avidity scores using both IFA and ELISA testing formats. **Note: Avidity testing will not replace confirmatory neutralization testing, non-WNV flavivirus IgG antibody (Eg. dengue, SLE, etc.) may bind to the antigen preparations used in avidity assays.**

### Confirmed Case Diagnostic Test Criteria:

It is currently recommended that health jurisdictions/authorities use the Confirmed Case Diagnostic Test Criteria to confirm index cases (locally acquired) in their area each year; for subsequent cases, health jurisdictions/authorities could use the Probable Case Diagnostic Test Criteria to classify cases in their area as “confirmed”, **for the purposes of surveillance**. Throughout the remainder of the transmission season health jurisdictions/authorities may wish to document PRNT antibody titres to West Nile virus in a proportion of cases, to be determined by that health jurisdiction/authority, in order to rule-out the possibility of concurrent activity by other flaviviruses. [For further information on diagnostic testing algorithms for West Nile virus, see the section entitled Laboratory Specimen Diagnostic Testing Algorithm in Appendix 4 of the National Guidelines for Response to West Nile virus.]

### AT LEAST ONE of the following:

A 4-fold or greater change in WN virus neutralizing antibody titres (using a PRNT or other kind of neutralization assay) in paired acute and convalescent sera, or CSF. <b>OR</b>
Isolation of WN virus from, or demonstration of WN virus antigen or WN virus-specific genomic sequences in tissue, blood, CSF or other body fluids <b>OR</b>
Demonstration of flavivirus antibodies in a single serum or CSF sample using a WN virus IgM ELISA <sup>8,9</sup> , confirmed by the detection of WN virus specific antibodies using a PRNT (acute or convalescent specimen). <b>OR</b>
A 4-fold or greater change in flavivirus HI titres in paired acute and convalescent sera or demonstration of a seroconversion using a WN virus IgG ELISA <sup>8,9</sup> <b>AND</b> the detection of WN specific antibodies using a PRNT (acute or convalescent serum sample).

<sup>8</sup> Both CDC and commercial IgM / IgG ELISAs are now available for front line serological testing. Refer to appropriate assay procedures and kit inserts for the interpretation of test results.

**Note:** WNV IgM antibody may persist for more than a year and the demonstration of IgM antibodies in a patient's serum, particularly in residents of endemic areas, may not be diagnostic of an *acute* WNV viral infection. Seroconversion (by HI, IgG ELISA or PRNT assays) demonstrates a current WNV infection. Therefore, the collection of acute and convalescent sera for serologic analysis is particularly important to rule out diagnostic misinterpretation early in the WNV season (e.g. May, June) and to identify initial cases in a specific jurisdiction. However, it should be noted that seroconversions may not always be documented due to timing of acute sample collection (i.e. titres in acute sera may have already peaked). If static titres are observed in acute and convalescent paired sera, it is still possible the case may represent a recent infection. To help resolve this, the use of IgG avidity testing<sup>9</sup> may be considered to distinguish between current and past infection. The presence of both IgM antibody and low avidity IgG in a patient's convalescent serum sample are consistent with current cases of viral associated illness. However test results that show the presence of IgM and high avidity IgG are indicative of exposures that have occurred in the previous season. Immunocompromised individuals may not be able to mount an immune response necessary for a serological diagnosis. West Nile virus diagnostic test criteria for these individuals should be discussed with a medical microbiologist.

<sup>9</sup> Early in infection the immune system generates antibodies that bind relatively weakly to viral antigen (low avidity). As the infection proceeds, an increasing percentage of newly generated IgG antibody displays higher binding affinity to virus antigen and thus avidity also rises (Note: avidity is usually measured based upon the ability of IgG to dissociate from antigen preparations after incubation with a solution of urea). As long as high avidity IgG is not yet detected in the serum it can be assumed that the individual was exposed to the viral agent during a recent exposure. With respect to WNV infection it has not been precisely determined when (i.e. post-exposure) high avidity antibodies reach levels in serum that can be accurately detected by serological assays (there may be significant variation depending on the individual). However, it has been shown that greater than 95% of sera collected from individuals exposed to WNV 6-8 months previously will have IgG antibodies that bind strongly to viral antigen and will give high avidity scores using both IFA and ELISA testing formats. **Note: Avidity testing will not replace confirmatory neutralization testing, non-WNV flavivirus IgG antibody (Eg. dengue, SLE, etc.) may bind to the antigen preparations used in avidity assays.**