

Ontario Drug Benefit Formulary/Comparative Drug Index

Edition 43

Summary of Changes – January 2023
Effective January 31, 2023

Drug Programs Policy and Strategy Branch
Health Programs and Delivery Division
Ministry of Health

[Visit Formulary Downloads: Edition 43](#)

Table of Contents

| | |
|---|----|
| New Single Source Products..... | 3 |
| New Multi-Source Products..... | 4 |
| New Off-Formulary Interchangeable (OFI) Products..... | 9 |
| Revision of Limited Use Criteria | 11 |
| Removal of Therapeutic Note..... | 27 |
| Product Brand Name Changes | 28 |
| Product Brand and Manufacturer Name Changes | 29 |
| Drug Benefit Price (DBP) Changes | 30 |
| Discontinued Products | 32 |
| Delisted Products | 33 |

New Single Source Products

Generic Name: INSULIN ASPART

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr | DBP |
|----------|------------|----------|---------------------------|-----|------------|
| 02520974 | Kirsty | 100U/mL | Inj Sol-5x3mL Pref Pen Pk | BGP | 42.7125/Pk |

New Multi-Source Products

Where applicable, please consult the respective brand reference product's drug profile on the ODB e-Formulary for the details of the Limited Use (LU) code and criteria, and/or any associated Therapeutic Notes (TN).

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-------------------------|----------|-------------|-----|--------|
| 02525348 | Amoxicillin Capsules BP | 250mg | Cap | SAI | 0.0672 |
| 02525356 | Amoxicillin Capsules BP | 500mg | Cap | SAI | 0.1308 |

(Interchangeable with Amoxil – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|----------------|----------|-------------|-----|--------|
| 02482630 | Apo-Ticagrelor | 90mg | Tab | APX | 0.3960 |

(Interchangeable with Brilinta – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|---------------|----------|-------------|-----|--------|
| 02486806 | Auro-Apixaban | 2.5mg | Tab | AUR | 0.4084 |

(Interchangeable with Eliquis DIN 02377233 – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|---------------|----------|-------------|-----|--------|
| 09858239 | Auro-Apixaban | 2.5mg | Tab | AUR | 0.4084 |

(Interchangeable with Eliquis PIN 09857463 – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|---------------|----------|-------------|-----|--------|
| 02486814 | Auro-Apixaban | 5mg | Tab | AUR | 0.4084 |

(Interchangeable with Eliquis – LU)

New Multi-Source Products (Continued)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|------------------|----------|-------------|-----|---------|
| 02530007 | Auro-Tofacitinib | 5mg | Tab | AUR | 5.9897 |
| 02530015 | Auro-Tofacitinib | 10mg | Tab | AUR | 21.1718 |

(Interchangeable with Xeljanz – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-------------------|----------|-------------|-----|--------|
| 02528037 | Jamp Diltiazem CD | 120mg | CD Cap | JPC | 0.3634 |
| 02528045 | Jamp Diltiazem CD | 180mg | CD Cap | JPC | 0.4824 |
| 02528053 | Jamp Diltiazem CD | 240mg | CD Cap | JPC | 0.6399 |
| 02528061 | Jamp Diltiazem CD | 300mg | CD Cap | JPC | 0.7999 |

(Interchangeable with Cardizem CD – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|----------------------|----------|-------------|-----|--------|
| 02245946 | Jamp-Docusate Sodium | 100mg | Cap | JPC | 0.0328 |

(Interchangeable with Colace – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|--|-------------------|-------------|-----|---------|
| 02519461 | Jamp Efavirenz/Emtricitabine/ Tenofovir Disoproxil Fumarate | 600mg/200mg/300mg | Tab | JPC | 11.3300 |

(Interchangeable with Atripla – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|----------------|----------|-------------|-----|---------|
| 02520354 | Jamp Linezolid | 600mg | Tab | JPC | 19.3041 |

(Interchangeable with Zyvoxam – LU)

New Multi-Source Products (Continued)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|---------------------------|----------|-------------|-----|--------|
| 02527200 | Jamp Perindopril Erbumine | 2mg | Tab | JPC | 0.1632 |
| 02527219 | Jamp Perindopril Erbumine | 4mg | Tab | JPC | 0.2042 |
| 02527227 | Jamp Perindopril Erbumine | 8mg | Tab | JPC | 0.2831 |

(Interchangeable with Coversyl – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|---------------|----------|-------------|-----|--------|
| 02495449 | Mint-Apixaban | 5mg | Tab | MIN | 0.4084 |

(Interchangeable with Eliquis – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|--------------------|----------|-------------|-----|--------|
| 02522225 | Mint-Quetiapine XR | 400mg | ER Tab | MIN | 1.3270 |

(Interchangeable with Seroquel XR – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-----------------|----------|-------------|-----|--------|
| 02526379 | Mint-Ranitidine | 150mg | Tab | MIN | 0.1197 |
| 02526387 | Mint-Ranitidine | 300mg | Tab | MIN | 0.2253 |

(Interchangeable with Zantac – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-----------------|----------|-------------|-----|--------|
| 02522101 | Nat-Montelukast | 4mg | Chew Tab | NAT | 0.2758 |

(Interchangeable with Singulair – LU)

New Multi-Source Products (Continued)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-----------------|----------|-------------|-----|--------|
| 02527014 | NRA-Candesartan | 8mg | Tab | NRA | 0.2281 |
| 02527022 | NRA-Candesartan | 16mg | Tab | NRA | 0.2281 |
| 02527030 | NRA-Candesartan | 32mg | Tab | NRA | 0.2281 |

(Interchangeable with Atacand – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|----------------|----------|-------------|-----|--------|
| 02477645 | NRA-Citalopram | 20mg | Tab | NRA | 0.1332 |
| 02477653 | NRA-Citalopram | 40mg | Tab | NRA | 0.1332 |

(Interchangeable with Celexa – GB)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-----------------|----------|-------------|-----|--------|
| 02522799 | PMS-Tofacitinib | 5mg | Tab | PMS | 5.9897 |

(Interchangeable with Xeljanz – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|------------------|----------|-------------|-----|---------|
| 02511304 | Taro-Tofacitinib | 5mg | Tab | TAR | 5.9897 |
| 02511312 | Taro-Tofacitinib | 10mg | Tab | TAR | 21.1718 |

(Interchangeable with Xeljanz – LU)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP |
|----------|-------------------|----------|-------------|-----|--------|
| 02429322 | Teva-Diltiazem XC | 180mg | ER Tab | TEV | 0.9195 |
| 02429330 | Teva-Diltiazem XC | 240mg | ER Tab | TEV | 1.2212 |
| 02429349 | Teva-Diltiazem XC | 300mg | ER Tab | TEV | 1.2175 |
| 02429357 | Teva-Diltiazem XC | 360mg | ER Tab | TEV | 1.2211 |

(Interchangeable with Tiazac XC – GB)

New Off-Formulary Interchangeable (OFI) Products

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|----------------|----------|-------------|-----|-----------|
| 02518910 | GLN-Apremilast | 30mg | Tab | GLP | 18.7239 |

(Interchangeable with Otezla)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|----------------|----------|-------------|-----|-----------|
| 02528207 | Jamp Clonidine | 0.025mg | Tab | JPC | 0.2713 |

(Interchangeable with Dixarit)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|--------------|----------|-------------|-----|-----------|
| 02530244 | Modafinil | 100mg | Tab | SAI | 0.9293 |

(Interchangeable with Alertec)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|--------------|----------|---------------------------------------|-----|-----------|
| 02529076 | Moxifloxacin | 0.5% w/v | Oph Sol-3mL Pk (Preservative-Free) | SAI | 11.2700 |

(Interchangeable with Vigamox)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|--------------|----------|-------------|-----|-----------|
| 02524511 | M-Valsartan | 40mg | Tab | MAT | 0.5823 |

(Interchangeable with Diovan)

New Off-Formulary Interchangeable (OFI) Products (Continued)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|-----------------|----------|-------------|-----|-----------|
| 02522128 | Nat-Montelukast | 5mg | Chew Tab | NAT | 1.2077 |
| 02522136 | Nat-Montelukast | 10mg | Tab | NAT | 1.7737 |

(Interchangeable with Singulair)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|---------------------|----------|-------------|-----|-----------|
| 02489384 | NRA-Rizatriptan ODT | 10mg | ODT | NRA | 11.1150 |

(Interchangeable with Maxalt RPD)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|------------------|----------|-------------|-----|-----------|
| 02489392 | NRA-Zolmitriptan | 2.5mg | Tab | NRA | 6.8600 |

(Interchangeable with Zomig)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|----------------|----------|-------------|-----|-----------|
| 02521733 | PMS-Apremilast | 30mg | Tab | PMS | 18.7238 |

(Interchangeable with Otezla)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | Unit Cost |
|----------|-------------------|----------|-------------|-----|-----------|
| 02529084 | Sandoz Apremilast | 30mg | Tab | SDZ | 18.7238 |

(Interchangeable with Otezla)

Revision of Limited Use Criteria

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|------------|------------|----------------|-----|
| 02419475 | Inflectra | 100mg/Vial | Inj Pd-Vial Pk | CEH |

Limited Use Codes Revised Clinical Criteria

Code 477

Ulcerative Colitis

For the treatment of moderate to severe ulcerative colitis in patients who meet the following criteria:

- A. Mayo score greater than or equal to 6 with an endoscopic subscore* of at least 2 (or other validated disease activity score confirming moderate to severe disease);
AND
- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine);
OR
Conventional treatment with a corticosteroid is contraindicated;
AND
- C. Infliximab is being used to induce remission or as a steroid-sparing maintenance therapy.

*The endoscopy procedure must be done within the 12 months prior to initiation of treatment.

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses may be considered in patients who have failed to respond to lower doses.)

Revision to Limited Use Criteria (Continued)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., Mayo score less than 6), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild ulcerative colitis (e.g., Mayo score less than 6) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Code 478

Luminal Crohn's disease

For the treatment of moderate to severe (luminal) Crohn's disease in patients who meet the following criteria:

- A. Harvey Bradshaw Index (HBI) score greater than or equal to 7 (or other validated disease activity score confirming moderate to severe disease);
AND
- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine, methotrexate);
OR
Conventional treatment with a corticosteroid is contraindicated;
AND

Revision to Limited Use Criteria (Continued)

- C. Infliximab is being used to induce remission or as a steroid-sparing maintenance therapy.

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses up to 10mg/kg/dose may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., HBI score decrease greater than or equal to 50% from pre-treatment measurement), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild Crohn's disease (e.g., HBI less than 7) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Revision to Limited Use Criteria (Continued) Code 479

Fistulising Crohn's disease

For the treatment of fistulising Crohn's disease in patients who meet the following criteria:

- Patient has actively draining perianal or enterocutaneous fistula(e) that have recurred OR persist despite a course of appropriate antibiotic therapy (e.g., ciprofloxacin and/or metronidazole)

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses up to 10mg/kg/dose may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and achieve and maintain response to therapy (e.g., partial or complete resolution of fistulae and symptom improvement).

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

LU Authorization Period: 1 year

Revision to Limited Use Criteria (Continued)

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|------------|------------|----------------|-----|
| 02470373 | Renflexis | 100mg/Vial | Inj Pd-Vial Pk | SAM |

Limited Use Codes Revised Clinical Criteria

Code 545

Ulcerative Colitis

For the treatment of moderate to severe ulcerative colitis in patients who meet the following criteria:

- A. Mayo score greater than or equal to 6 with an endoscopic subscore* of at least 2 (or other validated disease activity score confirming moderate to severe disease);
AND
- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine);
OR
Conventional treatment with a corticosteroid is contraindicated;
AND
- C. Infliximab is being used to induce remission or as a steroid-sparing maintenance therapy.

*The endoscopy procedure must be done within the 12 months prior to initiation of treatment.

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission.

Revision to Limited Use Criteria (Continued)

Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., Mayo score less than 6), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild ulcerative colitis (e.g., Mayo score less than 6) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Code 546

Luminal Crohn's disease

For the treatment of moderate to severe (luminal) Crohn's disease in patients who meet the following criteria:

- A. Harvey Bradshaw Index (HBI) score greater than or equal to 7 (or other validated disease activity score confirming moderate to severe disease);
AND
- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine, methotrexate);
OR
Conventional treatment with a corticosteroid is contraindicated;
AND
- C. Infliximab is being used to induce remission or as a steroid-sparing maintenance therapy.

Revision to Limited Use Criteria (Continued)

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses up to 10mg/kg/dose may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., HBI score decrease greater than or equal to 50% from pre-treatment measurement), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild Crohn's disease (e.g., HBI less than 7) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Code 547

Fistulising Crohn's disease

For the treatment of fistulising Crohn's disease in patients who meet the following criteria:

- Patient has actively draining perianal or enterocutaneous fistula(e) that have recurred OR persist despite a course of appropriate antibiotic therapy (e.g., ciprofloxacin and/or metronidazole)

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

Revision to Limited Use Criteria (Continued)

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses up to 10mg/kg/dose may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and achieve and maintain response to therapy (e.g., partial or complete resolution of fistulae and symptom improvement).

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

LU Authorization Period: 1 year

Revision to Limited Use Criteria (Continued)

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|-------------------|------------|--|-----|
| 02511045 | Abrilada | 40mg/0.8mL | Inj Sol-0.8mL Pref Pen (Preservative-Free) | PFI |
| 02511053 | Abrilada | 40mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | PFI |
| 02459310 | Amgevita | 20mg/0.4mL | Inj Sol-0.4mL Pref Syr (Preservative-Free) | AMG |
| 02459302 | Amgevita | 40mg/0.8mL | Inj Sol-0.8mL Pref Autoinj (Preservative-Free) | AMG |
| 02459299 | Amgevita | 40mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | AMG |
| 02473097 | Hadlima | 40mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | SAM |
| 02473100 | Hadlima PushTouch | 40mg/0.8mL | Inj Sol-0.8mL Pref Autoinj (Preservative-Free) | SAM |
| 02502380 | Hulio | 20mg/0.4mL | Inj Sol-0.4mL Pref Syr (Preservative-Free) | BGP |
| 02502402 | Hulio | 40mg/0.8mL | Inj Sol-0.8mL Pref Pen (Preservative-Free) | BGP |
| 02502399 | Hulio | 40mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | BGP |
| 02505258 | Hyrimoz | 20mg/0.4mL | Inj Sol-0.4mL Pref Syr (Preservative-Free) | SDZ |
| 02492156 | Hyrimoz | 40mg/0.8mL | Inj Sol-0.8mL Pref Autoinj (Preservative-Free) | SDZ |
| 02492164 | Hyrimoz | 40mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | SDZ |
| 02502674 | Idacio | 40mg/0.8mL | Inj Sol-0.8mL Pref Pen (Preservative-Free) | FKC |
| 02502682 | Idacio | 40mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | FKC |
| 02523949 | Simlandi | 40mg/0.4mL | Inj Sol-0.4mL Pref Syr (Preservative-Free) | JPC |
| 02523957 | Simlandi | 40mg/0.4mL | Inj Sol-0.4mL Pref Autoinj Syr (Preservative-Free) | JPC |
| 02523965 | Simlandi | 80mg/0.8mL | Inj Sol-0.8mL Pref Syr (Preservative-Free) | JPC |
| 02523779 | Yuflyma | 40mg/0.4mL | Inj Sol-0.4mL Pref Autoinj Pen (Preservative-Free) | CEH |
| 02523760 | Yuflyma | 40mg/0.4mL | Inj Sol-0.4mL Pref Syr (Preservative-Free) | CEH |

Revision to Limited Use Criteria (Continued)
Limited Use Codes Revised Clinical Criteria

Code 604

Luminal Crohn's disease

For the treatment of moderate to severe (luminal) Crohn's disease in patients who meet the following criteria:

- A. Harvey Bradshaw Index (HBI) score greater than or equal to 7 (or other validated disease activity score confirming moderate to severe disease);

AND

- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);

OR

Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine, methotrexate);

OR

Conventional treatment with a corticosteroid is contraindicated;

AND

- C. Adalimumab is being used to induce remission or as a steroid-sparing maintenance therapy.

The recommended induction dosing regimen is 160mg at week 0, followed by 80mg at week 2.

The recommended maintenance dosing regimen is up to 40mg every 2 weeks. (Note: higher doses may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., HBI score decrease greater than or equal to 50% from pre-treatment measurement), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Revision to Limited Use Criteria (Continued)

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild Crohn's disease (e.g., HBI less than 7) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Code 605

Fistulising Crohn's disease

For the treatment of fistulising Crohn's disease with concomitant luminal disease in patients who meet the following criteria:

- A. Patient has actively draining perianal or enterocutaneous fistula(e) that have recurred OR persist despite a course of appropriate antibiotic therapy (e.g., ciprofloxacin and/or metronidazole);
AND
- B. Harvey Bradshaw Index (HBI) score greater than or equal to 7 (or other validated disease activity score confirming moderate to severe disease)

The recommended induction dosing regimen is 160mg at week 0, followed by 80mg at week 2.

The recommended maintenance dosing regimen is up to 40mg every 2 weeks. (Note: higher doses may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and achieve and maintain response to therapy (e.g., partial or complete resolution of fistulae and symptom improvement).

Revision to Limited Use Criteria (Continued)

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

LU Authorization Period: 1 year

Code 606

Ulcerative Colitis

For the treatment of moderate to severe ulcerative colitis in patients who meet the following criteria:

- A. Mayo score greater than or equal to 6 with an endoscopic subscore* of at least 2 (or other validated disease activity score confirming moderate to severe disease);
AND
- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week)
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine)
OR
Conventional treatment with a corticosteroid is contraindicated;
AND
- C. Adalimumab is being used to induce remission or as a steroid-sparing maintenance therapy.

*The endoscopy procedure must be done within the 12 months prior to initiation of treatment.

The recommended induction dosing regimen is up to 160mg at week 0, followed by up to 80mg at week 2.

The recommended maintenance dosing regimen is up to 40mg every 2 weeks. (Note: higher doses may be considered in patients who have failed to respond to lower doses.)

Revision to Limited Use Criteria (Continued)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., Mayo score less than 6), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild ulcerative colitis (e.g., Mayo score less than 6) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|------------|------------|--------------------|-----|
| 02496933 | Avsola | 100mg/Vial | Pd for Sol-Vial Pk | AMG |

Limited Use Codes Revised Clinical Criteria

Code 596

Ulcerative Colitis

For the treatment of moderate to severe ulcerative colitis in patients who meet the following criteria:

- A. Mayo score greater than or equal to 6 with an endoscopic subscore* of at least 2 (or other validated disease activity score confirming moderate to severe disease);

AND

Revision to Limited Use Criteria (Continued)

- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine);
OR
Conventional treatment with a corticosteroid is contraindicated;
AND
C. Infliximab is being used to induce remission or as a steroid-sparing maintenance therapy.

*The endoscopy procedure must be done within the 12 months prior to initiation of treatment.

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., Mayo score less than 6), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild ulcerative colitis (e.g., Mayo score less than 6) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Revision to Limited Use Criteria (Continued)

Code 597

Luminal Crohn's disease

For the treatment of moderate to severe (luminal) Crohn's disease in patients who meet the following criteria:

- A. Harvey Bradshaw Index (HBI) score greater than or equal to 7 (or other validated disease activity score confirming moderate to severe disease);
AND
- B. Failed conventional treatment with a corticosteroid (prednisone 40–60mg/day [or equivalent]) for a minimum of 14 days (or intravenous corticosteroid for 1 week);
OR
Responded to/stabilized on conventional treatment with a corticosteroid, with or without an immunosuppressant (e.g., azathioprine, 6-mercaptopurine, methotrexate);
OR
Conventional treatment with a corticosteroid is contraindicated;
AND
- C. Infliximab is being used to induce remission or as a steroid-sparing maintenance therapy.

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses up to 10mg/kg/dose may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and have demonstrated a treatment response or are in remission. Examples of treatment response include clinically meaningful reductions in disease activity scores (e.g., HBI score decrease greater than or equal to 50% from pre-treatment measurement), along with improvements in endoscopic findings and reduction or discontinuation of corticosteroids.

Prescribers may wish to consider other funded alternatives for patients unable to discontinue corticosteroid therapy.

Revision to Limited Use Criteria (Continued)

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

Patients with mild Crohn's disease (e.g., HBI less than 7) may be considered on a case-by-case basis through the Exceptional Access Program.

LU Authorization Period: 1 year

Code 598

Fistulising Crohn's disease

For the treatment of fistulising Crohn's disease in patients who meet the following criteria:

- Patient has actively draining perianal or enterocutaneous fistula(e) that have recurred OR persist despite a course of appropriate antibiotic therapy (e.g., ciprofloxacin and/or metronidazole)

The recommended induction dosing regimen is 5mg/kg/dose at 0, 2, and 6 weeks.

The recommended maintenance dosing regimen is 5mg/kg/dose every 8 weeks. (Note: higher doses up to 10mg/kg/dose may be considered in patients who have failed to respond to lower doses.)

Maintenance/Renewal:

Maintenance therapy is funded for patients who met the initiation criteria and achieve and maintain response to therapy (e.g., partial or complete resolution of fistulae and symptom improvement).

Exclusion criteria (initial and renewal coverage):

- Combination therapy with another biologic used to treat inflammatory bowel disease will not be funded.

LU Authorization Period: 1 year

Removal of Therapeutic Note

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|--------------------|----------|-------------|-----|
| 09857268 | Mirapex | 0.25mg | Tab | BOE |
| 02292378 | Apo-Pramipexole | 0.25mg | Tab | APX |
| 02424061 | Auro-Pramipexole | 0.25mg | Tab | AUR |
| 02297302 | Co Pramipexole | 0.25mg | Tab | COB |
| 02367602 | Pramipexole | 0.25mg | Tab | SAI |
| 02309122 | Pramipexole | 0.25mg | Tab | SIV |
| 02315262 | Sandoz Pramipexole | 0.25mg | Tab | SDZ |

Therapeutic Note to be removed:

NOTE: Mirapex is indicated for both the symptomatic treatment of idiopathic Parkinson's Disease and moderate to severe idiopathic Restless Legs Syndrome under the manufacturer's Drug Identification Number (DIN). Mirapex has also been assigned a Product Identification Number (PIN) for the indication of Parkinson's Disease specifically. Apo-Pramipexole, Auro-Pramipexole, Novo-Pramipexole, Sandoz Pramipexole, Co Pramipexole and Mylan-Pramipexole products are interchangeable with Mirapex for the treatment of Parkinson's Disease.

All generic pramipexole 0.25mg Tab products are now in the same interchangeable category with Mirapex 0.25mg Tab DIN 02237145. The Mirapex PIN 09857268 is delisted.

Product Brand Name Changes

| DIN/PIN | Current Product Name | New Product Name | MFR | Strength | Dosage Form |
|----------|----------------------|------------------|-----|----------|-------------|
| 02296349 | Act Ondansetron | Teva-Ondansetron | TEV | 4mg | Tab |
| 02296357 | Act Ondansetron | Teva-Ondansetron | TEV | 8mg | Tab |

Product Brand and Manufacturer Name Changes

| DIN/PIN | Current Product Name | Current Mfr | New Product Name | New Mfr | Strength | Dosage Form |
|----------|----------------------|-------------|------------------|---------|----------|-------------|
| 02297302 | Co Pramipexole | COB | Act Pramipexole | TEV | 0.25mg | Tab |
| 02297310 | Co Pramipexole | COB | Act Pramipexole | TEV | 0.5mg | Tab |
| 02297329 | Co Pramipexole | COB | Act Pramipexole | TEV | 1mg | Tab |
| 02297337 | Co Pramipexole | COB | Act Pramipexole | TEV | 1.5mg | Tab |

Drug Benefit Price (DBP) Changes

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP/Unit Price |
|----------|------------------------|---------------------------------|------------------|-----|----------------|
| 02441934 | Act Methylphenidate ER | 18mg | SR Tab | TEV | 1.0493 |
| 02441942 | Act Methylphenidate ER | 27mg | SR Tab | TEV | 1.2109 |
| 02441950 | Act Methylphenidate ER | 36mg | SR Tab | TEV | 1.3726 |
| 02441969 | Act Methylphenidate ER | 54mg | SR Tab | TEV | 1.6958 |
| 02244394 | Apo-Cefuroxime | 500mg | Tab | APX | 1.6616 |
| 02452731 | Apo-Methylphenidate ER | 18mg | SR Tab | APX | 1.0493 |
| 02452758 | Apo-Methylphenidate ER | 27mg | SR Tab | APX | 1.2109 |
| 02452766 | Apo-Methylphenidate ER | 36mg | SR Tab | APX | 1.3726 |
| 02330377 | Apo-Methylphenidate ER | 54mg | SR Tab | APX | 1.6958 |
| 02426552 | Apo-Linezolid | 600mg | Tab | APX | 19.3041 |
| 02344831 | Auro-Cefuroxime | 500mg | Tab | APX | 1.6616 |
| 02247732 | Concerta | 18mg | SR Tab | JAN | 3.0171 |
| 02250241 | Concerta | 27mg | SR Tab | JAN | 3.4819 |
| 02247733 | Concerta | 36mg | SR Tab | JAN | 3.9468 |
| 02247734 | Concerta | 54mg | SR Tab | JAN | 4.8761 |
| 00263818 | Cotazym | 8000 & 30000 & 3000 USP Units | Cap | ORG | 0.2520 |
| 00502790 | Cotazym ECS 8 | 8000 & 30000 & 3000 USP Units | Ent Microsph Cap | ORG | 0.4549 |
| 00821373 | Cotazym ECS 20 | 20000 & 55000 & 55000 USP Units | Ent Microsph Cap | ORG | 1.1928 |
| 02298791 | Emend | 80mg | Cap | MEK | 35.3082 |
| 02298805 | Emend | 125mg | Cap | MEK | 34.9776 |
| 02298813 | Emend Tri-Pack | 125mg & 80mg | Cap | MEK | 105.9246 |

Drug Benefit Price (DBP) Changes (Continued)

| DIN/PIN | Product Name | Strength | Dosage Form | Mfr | DBP/Unit Price |
|----------|------------------|----------------------------|---------------------------|-----|----------------|
| 02042487 | Marvelon 21 | 0.15mg & 0.03mg | Tab-21 Pk | ORG | 20.1010 |
| 02042479 | Marvelon 28 | 0.15mg & 0.03mg | Tab-28 Pk | ORG | 20.1010 |
| 02240521 | Maxalt | 10mg | Tab | OCI | 21.4865 |
| 02240518 | Maxalt RPD | 5mg | Orally Disintegrating Tab | OCI | 21.4865 |
| 02240519 | Maxalt RPD | 10mg | Orally Disintegrating Tab | OCI | 21.4865 |
| 02529769 | M-Ticagrelor | 90mg | Tab | MAT | 0.3960 |
| 02243796 | Pariet | 10mg | Tab | JAN | 1.4651 |
| 02243797 | Pariet | 20mg | Tab | JAN | 2.9301 |
| 02236950 | Risperdal | 1mg/mL | O/L | JAN | 1.5430/mL |
| 02255707 | Risperdal Consta | 25mg | Pd for Inj-Vial Pk | JAN | 191.1500 |
| 02255723 | Risperdal Consta | 37.5mg | Pd for Inj-Vial Pk | JAN | 286.7100 |
| 02255758 | Risperdal Consta | 50mg | Pd for Inj-Vial Pk | JAN | 382.2800 |
| 02422689 | Sandoz Linezolid | 600mg | Tab | SDZ | 19.3041 |
| 02243602 | Singulair | 4mg | Chew Tab | OCI | 1.5982 |
| 02231347 | Sporanox | 10mg/mL | Oral Sol | JAN | 1.1164/mL |
| 02047454 | Sporanox | 100mg | Cap | JAN | 5.9810 |
| 02492598 | Taro-Ticagrelor | 90mg | Tab | TAR | 0.3960 |
| 02230893 | Topamax | 25mg | Tab | JNO | 2.0797 |
| 02230894 | Topamax | 100mg | Tab | JNO | 3.9000 |
| 02230896 | Topamax | 200mg | Tab | JNO | 5.7585 |
| 02239907 | Topamax Sprinkle | 15mg | Sprinkle Cap | JNO | 1.9800 |
| 02239908 | Topamax Sprinkle | 25mg | Sprinkle Cap | JNO | 2.0700 |
| 02244981 | Tracleer | 62.5mg | Tab | JAN | 78.4500 |
| 02244982 | Tracleer | 125mg | Tab | JAN | 78.4500 |
| 00556734 | Vermox | 100mg | Tab | JAN | 9.2650 |
| 02250519 | Zavesca | 100mg | Cap | ACT | 128.9600 |
| 02361752 | Zenhale | 100mcg & 5mcg Metered Dose | Inh-120 Dose Pk | OCI | 111.3000 |
| 02361760 | Zenhale | 200mcg & 5mcg Metered Dose | Inh-120 Dose Pk | OCI | 134.8600 |

Discontinued Products

(Some products will remain on Formulary for six months to facilitate depletion of supply)

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|------------------------------|----------|----------------------------------|-----|
| 00585092 | Depo-Provera | 150mg/mL | Inj | PFI |
| 01931563 | Gastrolyte | | Oral Pd-1 Sach Pk | SAV |
| 09858001 | InspiraChamber | | | INS |
| 09858002 | InspiraChamber + Mask Small | | | INS |
| 09858003 | InspiraChamber + Mask Medium | | | INS |
| 09858004 | InspiraChamber + Mask Large | | | INS |
| 02513447 | Riabni | 10mg/mL | Inj Sol-Vial (Preservative-Free) | AMG |
| 02353040 | Ropinirole | 0.25mg | Tab | SAI |
| 02353059 | Ropinirole | 1mg | Tab | SAI |
| 02303396 | Sandoz Metoprolol SR | 100mg | LA Tab | SDZ |
| 02303418 | Sandoz Metoprolol SR | 200mg | LA Tab | SDZ |

Delisted Products

| DIN/PIN | Brand Name | Strength | Dosage Form | Mfr |
|----------|------------------------|----------|-------------|-----|
| 02413736 | PMS-Methylphenidate ER | 27mg | SR Tab | PMS |
| 02413744 | PMS-Methylphenidate ER | 36mg | SR Tab | PMS |
| 00021261 | Teva-Chloroquine | 250mg | Tab | TEV |

